



ISSN: 2395-7476
IJHS 2016; 2(3): 123-126
© 2016 IJHS
www.homesciencejournal.com
Received: 21-07-2016
Accepted: 22-08-2016

Dr. Ekta Singh Chauhan
Department of Food Science and
Nutrition, Banasthali University,
Banasthali, Distt-Tonk. Rajasthan,
304022, India

Anamika Tiwari
Department of Food Science and
Nutrition, Banasthali
University, Banasthali, Distt-
Tonk. Rajasthan, 304022, India

Akriti Singh
Department of Food Science and
Nutrition, Banasthali University,
Banasthali, Distt-Tonk. Rajasthan,
304022, India

Correspondence
Anamika Tiwari
Department of Food Science and
Nutrition, Banasthali University,
Banasthali, Distt-Tonk. Rajasthan,
304022, India

Phytochemical screening of Knol-Khol (*Brassica caulorapa*) Powder and Juice - A comparative study

Dr. Ekta Singh Chauhan, Anamika Tiwari and Akriti Singh

Abstract

Cruciferous vegetables are one of the dominant food crops worldwide. Cruciferous vegetables are vegetables of family *Brassicaceae* (also called cruciferae). They are good source of many health promoting and potentially protective phytochemicals including folic acid, phenolics, carotenoids, selenium, glucosinolates and vitamin c. Incorporating these potent plant based compounds in daily food is a safe, effective and easily available way to guard against many of today's most common diseases. The Knol- Khol (*Brassica caulorapa*) is cool season cruciferous vegetable. Present study was conducted to assess the qualitative phytochemical analysis of knol-khol powder and juice. The extract of powder and juice of knol-khol reveals the presence of many phytochemicals such as alkaloids, glycosides, flavonoids, saponnin, tannin, steroids, terpenes and phytosterols. The compounds isolated from this vegetable possess very important biological activities including anti-diabetic, antioxidant, hypolipidemic, antihyperglycemic, Cardioprotective, and Anti-cancer.

Keywords: Flavonoids, saponnin, tannin, steroids, phytosterols, anti-diabetic and antihyperglycemic

Introduction

Cruciferous vegetables are “vegetables of the *Brassicaceae* family also called as cruciferae”. Generally consumed cruciferous vegetables are broccoli, brussels sprouts, kale, mustard, cabbage, turnips, cauliflower, boy choy and Chinese cabbage (Jane V. Higdon *et al.* 2007) [5]. Almost all parts of some species are used as food, including stem (kohlrabi), root (turnips, rutabaga), leaves (cabbage, collard greens), seeds (mustard seed and oil- producing rapeseed) and flower (broccoli, cauliflower). “Cruciferous vegetables are one of the prevailing food crops worldwide”. Brassica vegetables are greatly “regarded for their nutritional value, they are rich source of vit.c, soluble fiber as well as contain multiple nutrients and phytochemicals”. Phytochemicals are the compounds derived from plants hypothesized to be responsible for much of the disease protection in our body they are present in diet high in fruits, vegetables, cereals and plant based beverages (Arts, I.C. *et al.* 2005) [1]. Recent studies reveal that cruciferous are the good source of natural antioxidants because they “contain carotenoids, tocopherols and ascorbic acid”. Epidemiological studies show these compounds may help to protect the human body against free radicals. Principal are their antioxidants effects, which are responsible for scavenging free radicals from our body and prevent body against its harmful effects (Sharique Ahmed and Seerat Hussain Beigh, 2009) [9].

The cruciferous vegetables have glucosinolates and also contain myrosinase an enzyme in a different area of the cell. Myrosinase hydrolyzes glucosinolates, into secondary metabolites, such as isothiocyanates and indoles, which are considered to be responsible for the anticarcinogenic action of cruciferous vegetables of cruciferous vegetables. Brassica vegetables contain “indole-3-carbinol, a chemical which helps in DNA repair in cells and helps to obstruct the growth of cancer cells” (Fan *et al.* 2006 and Wu *et al.* 2010) [2, 11]. These vegetables are also a good source of carotenoids with broccoli containing higher level of carotenoids (Farnham *et al.* 2009) [3]. “Brassica vegetables are a potent modulator of the innate immune response system with potent antiviral, antibacterial and anticancer activity, (Vivar *et al.* 2009) [10] however, it also is an antiandrogen”. (Le *et al.* 2003) [7].

Knol-Khol also belongs to *Brassicaceae* family. Knol- Khol is biologically known as *Brassica oleracea var. gangylodes*.

It is also a cool season cruciferous vegetable. It is sweet and juicy and not a fruit but a swollen stem growing about the ground. Knol-Khol is locally known as Ganth ghobi or Kholrabi. In India kohlrabi is more commonly called Knol-Khol (English) or Nookal (Hindi). Knol- Khol (*Brassica oleracea*) is a perennial plant and is grown throughout the year. It has a very cool and juicy stem, which can be consumed. It is pale green or cranberry colour. Knol-Khol is rich in vitamin C, contains negligible amount of fats and calories. "It is rich in the antioxidants like vitamins C, E and carotene and is a good source of dietary fibers". Phytochemicals which are present in knol-khol are isothiocyanates, sulforaphane, and indol-3-carbinol.

Antioxidants which are present in Knol-Khol help to kill harmful radicals in our body, thus helps to fight against cancer. It is specially used in curing colon cancers. Dietary fibers present in Knol- Khol helps in effective functioning of our digestive system. They also contain bioactive substances like sulphoraphanes and isothiocyanates, which help in the production of protective enzymes in our body and make our immune system stronger (Kurilich A.C. *et al.* 2002) [6]. Sulforaphane belongs to a group of phytochemicals or disease fighting compounds in plants food, known as isothiocyanates. Along with related phytochemicals, it helps to prevent certain enzymes from activating cancer causing agents in body and increases the body production of enzymes that clean carcinogens out of the system before they can damage the cells.

Vijaykumar P. Rasal M. *et al.* (2006) [8] have concluded in his study that Knol-khol extract prepared in petroleum ether rejuvenate the pancreas as well as helps in lowering hyperglycemia and oxidative stress. "Aqueous extract of Knol-Khol has showed antidiabetic activity in alloxan induced diabetic rats". Therefore, "knol-khol is beneficial for the treatment of diabetes and many other associated diseases".

Materials and Methods

Plant material

Knol-khol leaves were shade dried followed by hot air oven drying at 50° centigrade and then ground to a fine powder and stored in air tight container for the analysis. Fresh knol-khol leaves were grinded in the mixer for the collection of juice.

Qualitative phytochemical evaluation- (Handa, 1995) [4]

Extraction

The coarse powder and juice of knol-khol was extracted with petroleum ether, chloroform, methanol and water at the ratio of 30:70. The extracts of knol-khol powder and juice were collected separately and filtered using Whatman filter paper. All the extracts were concentrated and the excessive solvents were evaporated under vacuum.

Preliminary phytochemical analysis

All plant extracts were further used for chemical tests for the presence of following phytochemicals such as phenolics compounds, alkaloids, saponin, glycosides, phytosterols, tannin, flavonoids, steroids terpenoids using the methods mentioned below:-

A. Alkaloids

a. Mayer's test

To a few ml of filtrate, a drop or two of Mayer's reagent were reagent were added by the side of test tube. A white or creamy precipitate indicate indicated the test as positive.

b. Wagner's test

To a few ml of filtrate, few drops of wagner's reagent were added by the side of the test tube. A reddish –brown precipitate confirmed the test as positive.

B. Glycosides

a. To ml of aqueous extract of the samples, 5ml of Benedict's solution and few drop of dilute HCl were added and heated for minutes. The solution became red with precipitate which indicated the presence of glycosides.

b. Brontrager's Test

To 2 ml of filtered hydrolysate, 3 ml of chloroform was added and shaken, chloroform layer was separated and 10% ammonia solution was added to it pink colour indicated the presence of glycosides.

C. Terpenoids

Liebermann – Burchard's test: 2ml of acetic anhydride solution was added to 1ml of petroleum ether extract of the drug in chloroform, followed by 1 ml of concentrated sulphuric acid. A violet color ring was formed indicating the presence of terpenoids.

D. Steroids

Liebermann –Burchard's test: 2 ml of acetic anhydride solution was added to 1 ml of petroleum ether extract of the drug in chloroform followed by 1 ml of concentrated sulphuric acid. A greenish color was developed which turned to blue.

E. Saponins

In a test containing about 5 ml of an aqueous extract of the drug, a drop of sodium bicarbonate solution was added. The mixture was shaken vigorously and left for 3 minutes. Honeycomb like froth was formed.

F. Tannins

To 1-2 ml of plant extract, a few drops of 5% FeCL₃ solution were added. A green color indicated the presence of gallotannins which brown color indicated tannins.

G. Phytosterol

a. Liebermann-buchard's test

The extract (50 mg) was dissolved in 2ml acetic acid anhydride. To this, one or two drops of concentrated sulphuric acid were added slowly along the side of the test tube. An array of color changes showed the presence of phytosterols.

b. The extract was treated with Salkowski's reagent

The yellowish colour with green fluorescence appearance indicated the presence of phytosterol in it.

H. Flavonoids

SHONODA TEST: In a test tube containing 0.5 ml of alcoholic extract of the drug, 5-10 drops of dilute HCL was added followed by small pieces of magnesium. In the presence of flavonoids, a reddish pink or brown colour produced.

Result and Discussion

In the present study the comparison between the phytochemical property of knol-khol powder and juice were estimated. The preliminary phytochemical investigation on knol-khol powder and juice extracts revealed the presence of various secondary metabolites such as alkaloids, glycosides, steroids, flavonoids, saponnin, tannin, terpenoids and phytosterols in the different extracts (Table 1 &2).

Table 1: Preliminary Phytochemical Screening of Knol-Khol Powder Extract.

Name of the chemical test	Petroleum ether extract	Chloroform extract	Methanol extract	Distill water extract
Alkaloids	-	-	+	+
Glycosides	-	+	-	+
Steroids	+	+	+	+
Flavonoids	+	-	+	+
Saponin	+	+	-	+
Tannin	-	-	+	+
Terpenoids	+	-	-	+
Phytosterols	-	-	+	+

The phytochemical analysis of Knol-khol powder showed the presence of various phytochemical compounds in powder extract. The phytochemicals such as alkaloid, glycosides, steroids, flavonoids, saponin, tannin, terpenoids and

phytosterols are present in distilled water extract where as in petroleum ether, chloroform and methanol extract some phytochemicals were absent.

Table 2: Preliminary Phytochemical Screening of Knol-Khol Juice Extract

Name of the chemical test	Petroleum ether extract	Chloroform extract	Methanol extract	Distill water extract
Alkaloids	+	+	+	+
Glycosides	+	-	+	+
Steroids	+	+	-	+
Flavonoids	+	-	+	+
Saponin	-	-	-	+
Tannin	+	+	-	+
Terpenoids	+	+	-	+
Phytosterols	-	-	+	+

In the water extract of Knol-khol juice showed the presences of all the phytochemicals, where as some phytochemical compound were absent in other extract Knol-khol juice. The estimation of phytochemical compounds in powder and juice extract showed that the water extract of powder and juice of Knol-khol were rich in phytochemical compound which may help in treating many disease. In other prepared extract some phytochemicals were absent which may due to presence of chemical in the extract which react with the compound and makes it invisible. And, thus the powder had lost their phytochemical compound which makes them weak to enhance the quality of bark and seed powder. Thus, the study revealed that the Knol-khol powder and juice have the potential to cure many disease.

Conclusion

From the ancient times, plants have been used for treatment of variety of disease. Thus, the present study revealed that a number of positive effects of Knol-khol such as phytochemicals were found which is beneficial for the health. The phytochemical such as alkaloids, glycosides, steroids, flavonoids, saponin, tannin, terpenoids and phytosterols were present which increases the medicinal potential of Knol-khol and thus can be used for the treatment of various diseases. Therefore, modern medicine has many side effects and it is not quite safer for human consumption. So it is better to adopt natural food which has no side effects and quite safer too for human consumption.

Acknowledgment

The authors are thankful to Prof. Aditya Shastri Vice-chancellor, Banasthali Vidhyapith, Rajasthan, India and our department of Home Science for providing us lab facilities and financial assistance.

References

- Arts IC, Hollma PC. Polyphenols and disease risk in epidemiologic studies. *American Journal of Clinical Nutrition*. 2005; 81(1):S317-S325.
- Fan S, Meng Q, Auborn K, Carter T, Rosen EM. BRCA1 and BRCA2 as molecular target for phytochemicals indole-3-carbinol and genistein in breast and prostate cancer cells. *British Journal of Cancer*. 2006; 94(3):407-426.
- Farnham, Mark W, Kopsell, Dean A. Importance of Genotype on Carotenoid and Chlorophyll Levels in Broccoli Heads. *Hortscience*. 2009; 44(5):1248-1253.
- Handa SS. Quality control and standardization of herbal material and traditional remedies. *East pharma*. 1995; 38:23-25.
- Jane, Higdon V, Barbara Delage, David Williams E, Roderick H. Dashwood. Cruciferous Vegetables and human cancer risk: Epidemiologic evidence and mechanistic basis. *Pharmacol Res*. 2007; 55(3):224-236.
- Kurilich AC, Jeffery EH, Juvik JA, Wallig MA, Klein PL. Antioxidant capacity of different broccoli (*Brassica oleracea*) genotypes using the oxygen radical absorbance capacity assay. *Journal of Agriculture and Food Chemistry*. 2002; 50(18):5053-5057.
- Le Hen T, Schaldach, Charlene M, Firestone Gary L, Bjeldanes, Leonard F. Plant –derived 3,3'-Diindolymethane is a Strong Androgen Antagonist in Human Prostate cancer Cells. *Journal of Biological Chemical*. 2003; 278(23):21136-21145.
- Rasal VP, Shetty BB, Sinnathambi A, Yeshmania S, Ashok P. Antihyperglycaemic and Antioxidant activity of Brassica Oleracea in Streptozotocin Diabetic Rats. *The Internet Journal of Pharmacology*, 2006, 4(2). DOI: 10.5580/1219
- Sharique Ahmed, Seerat Hussain Beigh. Ascorbic acid, carotenoids, total phenolic content and antioxidant activity of various genotypes of Brassica Oleracea encephala. *Journal of Medical and Biological science*. 2009; 3(1):1-8.
- Vivar Omar I, Lin Chia-Lie, Firestone Gary L, Bjeldanes, Leonard F. 3,3'-Diindolymethane induces a G1 arrest in human prostate cancer cells irrespective of androgen receptor and p53 status. *Biochemical Pharmacology*. 2009; 78(5):469-476.

11. Wu Yongsheng, Feng Xiaolin, jin Yucui, Wu Zhaojia, Hankey, William Paisie, Carolyn Li Lei, Fengjuan *et al.* A Novel Mechanism of Indole-3-carbinol Effects on Breast Carcinogenesis Involves Induction of cdc25A Degradation. *Cancer Prevention Research.* 2010; 3(7):818-828.